



## NRC Publications Archive Archives des publications du CNRC

### **Stable high volumetric production of glycosylated human recombinant IFN $\alpha$ 2b in HEK293 cells**

Loignon, Martin; Perret, Sylvie; Kelly, John; Boulais, Denise; Cass, Brian; Bisson, Louis; Afkhamizarreh, Fatemeh; Durocher, Yves

This publication could be one of several versions: author's original, accepted manuscript or the publisher's version. / La version de cette publication peut être l'une des suivantes : la version prépublication de l'auteur, la version acceptée du manuscrit ou la version de l'éditeur.

For the publisher's version, please access the DOI link below. / Pour consulter la version de l'éditeur, utilisez le lien DOI ci-dessous.

#### **Publisher's version / Version de l'éditeur:**

<https://doi.org/10.1186/1472-6750-8-65>

*BMC Biotechnology*, 8, p. 65, 2008

#### **NRC Publications Record / Notice d'Archives des publications de CNRC:**

<https://nrc-publications.canada.ca/eng/view/object/?id=56439eaf-1097-4e7f-8b39-6413af53fc3c>

<https://publications-cnrc.canada.ca/fra/voir/objet/?id=56439eaf-1097-4e7f-8b39-6413af53fc3c>

Access and use of this website and the material on it are subject to the Terms and Conditions set forth at

<https://nrc-publications.canada.ca/eng/copyright>

READ THESE TERMS AND CONDITIONS CAREFULLY BEFORE USING THIS WEBSITE.

L'accès à ce site Web et l'utilisation de son contenu sont assujettis aux conditions présentées dans le site

<https://publications-cnrc.canada.ca/fra/droits>

LISEZ CES CONDITIONS ATTENTIVEMENT AVANT D'UTILISER CE SITE WEB.

#### **Questions?** Contact the NRC Publications Archive team at

PublicationsArchive-ArchivesPublications@nrc-cnrc.gc.ca. If you wish to email the authors directly, please see the first page of the publication for their contact information.

**Vous avez des questions?** Nous pouvons vous aider. Pour communiquer directement avec un auteur, consultez la première page de la revue dans laquelle son article a été publié afin de trouver ses coordonnées. Si vous n'arrivez pas à les repérer, communiquez avec nous à PublicationsArchive-ArchivesPublications@nrc-cnrc.gc.ca.



## Research article

## Open Access

# Stable high volumetric production of glycosylated human recombinant IFN $\alpha$ 2b in HEK293 cells

Martin Loignon<sup>1</sup>, Sylvie Perret<sup>1</sup>, John Kelly<sup>2</sup>, Denise Boulais<sup>1</sup>, Brian Cass<sup>1</sup>, Louis Bisson<sup>1</sup>, Fatemeh Afkhamizarreh<sup>1,3</sup> and Yves Durocher\*<sup>1</sup>

Address: <sup>1</sup>National Research Council Canada, Animal Cell Technology Group, Bioprocess Sector, Biotechnology Research Institute, 6100 Royalmount Ave., Montreal, QC, H4P 2R2, Canada, <sup>2</sup>Institute for Biological Sciences 100 Sussex Drive, Room 3100 Ottawa, ON K1A 0R6, Canada and <sup>3</sup>Department of Biomedical Engineering, Faculty of Medicine, McGill University, 3775 University Street, Montreal, QC H3A 2B4, Canada

Email: Martin Loignon - martin.loignon@nrc.gc.ca; Sylvie Perret - sylvie.perret@nrc.gc.ca; John Kelly - john.kelly@nrc.gc.ca; Denise Boulais - boudaisdlm@hotmail.com; Brian Cass - brian.cass@nrc.gc.ca; Louis Bisson - louis.bisson@nrc.gc.ca; Fatemeh Afkhamizarreh - Fatemeh.Afkhamizzareh@nrc.gc.ca; Yves Durocher\* - yves.durocher@nrc.gc.ca

\* Corresponding author

Published: 27 August 2008

Received: 22 December 2007

BMC Biotechnology 2008, 8:65 doi:10.1186/1472-6750-8-65

Accepted: 27 August 2008

This article is available from: <http://www.biomedcentral.com/1472-6750/8/65>

© 2008 Loignon et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Abstract

**Background:** Mammalian cells are becoming the prevailing expression system for the production of recombinant proteins because of their capacity for proper protein folding, assembly, and post-translational modifications. These systems currently allow high volumetric production of monoclonal recombinant antibodies in the range of grams per litre. However their use for large-scale expression of cytokines typically results in much lower volumetric productivity.

**Results:** We have engineered a HEK293 cell clone for high level production of human recombinant glycosylated IFN $\alpha$ 2b and developed a rapid and efficient method for its purification. This clone steadily produces more than 200 mg (up to 333 mg) of human recombinant IFN $\alpha$ 2b per liter of serum-free culture, which can be purified by a single-step cation-exchange chromatography following media acidification and clarification. This rapid procedure yields 98% pure IFN $\alpha$ 2b with a recovery greater than 70%. Purified IFN $\alpha$ 2b migrates on SDS-PAGE as two species, a major 21 kDa band and a minor 19 kDa band. N-terminal sequences of both forms are identical and correspond to the expected mature protein. Purified IFN $\alpha$ 2b elutes at neutral pH as a single peak with an apparent molecular weight of 44,000 Da as determined by size-exclusion chromatography. The presence of intramolecular and absence of intermolecular disulfide bridges is evidenced by the fact that non-reduced IFN $\alpha$ 2b has a greater electrophoretic mobility than the reduced form. Treatment of purified IFN $\alpha$ 2b with neuraminidase followed by O-glycosidase both increases electrophoretic mobility, indicating the presence of sialylated O-linked glycan. A detailed analysis of glycosylation by mass spectroscopy identifies disialylated and monosialylated forms as the major constituents of purified IFN $\alpha$ 2b. Electron transfer dissociation (ETD) shows that the glycans are linked to the expected threonine at position 106. Other minor glycosylated forms and non-sialylated species are also detected, similar to IFN $\alpha$ 2b produced naturally by lymphocytes. Further, the HEK293-produced IFN $\alpha$ 2b is biologically active as shown with reporter gene and antiviral assays.

**Conclusion:** These results show that the HEK293 cell line is an efficient and valuable host for the production of biologically active and glycosylated human IFN $\alpha$ 2b.

## Background

Interferons (IFNs) are cytokines with major therapeutic applications based on their antiviral, antiproliferative, and immunomodulatory activities. Type I IFNs (IFN $\alpha/\beta$ ) are massively produced in most cell types in response to viral and other microbial infections, and play a vital role in innate resistance to a wide variety of viruses [1]. The IFN $\alpha$ 2 locus comprises three variants, IFN $\alpha$ 2a, IFN $\alpha$ 2b and IFN $\alpha$ 2c, IFN $\alpha$ 2b being the predominant one detected in human genomic DNA [2,3]. Some of the many diseases treated with IFN $\alpha$ 2b, alone or in combination, include type B [4] and C hepatitis [5], several cancers such as melanoma [6-8], Kaposi's sarcoma [9], chronic myeloid lymphoma [10,11], and angioblastoma [12]. In the particular case of hepatitis C, a disease affecting over 170 million individuals worldwide, the combination of IFN $\alpha$  and the viral inhibitor ribavirin has become the standard treatment [13-15]. The rising incidence of certain cancers and viral hepatitis [16,17], in addition to ongoing investigations of novel therapeutic applications [18] are increasing the needs for human recombinant IFN $\alpha$ 2b.

Human recombinant IFN $\alpha$ 2b used in the clinic is synthesized in bacterial systems. When *E. coli* are grown in optimal conditions, a few grams (3 to 5) of recombinant human IFN $\alpha$  per litre of culture can be produced [19-21]. Bacterially produced recombinant human IFN $\alpha$ 2b is misfolded and therefore requires refolding into its native conformation. Once purified and refolded, the recoveries are typically lower than 20% [19,20]. This refolding process also often results in loss of specific activity. In addition, bacterially produced recombinant human IFN $\alpha$ 2b lacks the post-translational O-glycosylation present on the naturally synthesized protein. This non-glycosylated form of human recombinant IFN $\alpha$ 2b has a shorter serum half-life than the glycosylated form [22]. The chemical conjugation of polyethylene glycol (PEG) molecules to the core peptide (pegylation) has improved the pharmacodynamics and pharmacokinetics of IFN $\alpha$ 2b by increasing the serum half-life [23]. However, the pegylation of IFN $\alpha$ 2b has been reported in some cases to reduce its biological activity [24]. It has also been shown that the size of PEG molecules and sites of attachment differentially interfere with the interaction and binding of IFN $\alpha$ 2b to its receptor [25]. Nevertheless, the US Food and Drug Administration approved PEG-IFN $\alpha$ 2b in 2001 and PEG-IFN $\alpha$ 2a in 2002 for the treatment of chronic hepatitis C virus infection. Another common problem associated with the use of refolded and pegylated IFN $\alpha$  (PEG-IFN $\alpha$ ) is the formation of neutralizing antibodies. Antibody formation against PEG-IFN $\alpha$  in HCV-infected patients has been associated with treatment failure [26,27]. In mice, the presence of contaminating partially unfolded IFN species appears to play a key role in the appearance of these antibodies [28]

Human and other mammalian cells are expression systems of choice for the production of secreted recombinant proteins such as antibodies, sometimes yielding up to hundreds of milligram to gram quantities of purified product per liter of culture [29-31]. However, the volumetric productivity of human cells for given proteins such as cytokines (i.e. IFN $\alpha$ 2b) is often lower by several orders of magnitude. Originally, IFN $\alpha$  for therapeutic use was purified from the human lymphoblastoid Namalwa cell line following induction with Sendai virus. Despite the production of an IFN $\alpha$  with high biological activity, Namalwa cells were abandoned due to a limited productivity unable to satisfy an ever-growing demand. Other systems have been tested for the production of IFN $\alpha$ 2b. For example, avian eggs have been used for the production of human recombinant IFN $\alpha$ 2b [32,33], although the glycosylation pattern significantly differs from IFN $\alpha$ 2b produced by human peripheral blood leucocytes. Glycosylated IFN $\alpha$ 2b can also be produced in insect cells, but glycosylation is of the potentially immunogenic high-mannose type and lacks sialylation [34]. These limitations suggest that mammalian cells are preferable hosts for the production of fully glycosylated IFN $\alpha$ 2b. Chinese hamster ovary (CHO) cells have been used for the production of various human recombinant interferons. Glycosylated and biologically active mouse IFN $\alpha$  [35] can be produced in CHO cells. Similarly, Rossman *et al* have reported the production of 120  $\mu$ g/mL of IFN $\alpha$ 2b using a glutamine synthase-amplified vector in the mouse myeloma cell line NS0 [36]. This is the highest level of glycosylated recombinant human IFN $\alpha$ 2b produced in a mammalian system reported to date. *In vitro*, the biological activity of NS0-produced IFN $\alpha$ 2b is very similar to that produced by Namalwa cells.


Here we have successfully engineered a non-amplified IFN-producing clone derived from the HEK293 mammalian cell line that produces hundreds of milligrams of IFN $\alpha$ 2b per liter of serum-free media. The volumetric production of IFN $\alpha$ 2b reproducibly exceeds 200 mg/L in batch culture and remains stable in the absence of selection for more than four months in culture. The purified IFN $\alpha$ 2b is glycosylated and biologically active. Together these results demonstrate that cost-effective production and purification of glycosylated IFN $\alpha$ 2b from human cells can be achieved.

## Results

### **Generation of a stable IFN $\alpha$ 2b-expressing HEK293 cell clone and production in fed-batch cultures**

The expression plasmid pYD7 encoding the human IFN $\alpha$ 2b gene codon-optimized for expression in human cells (Fig. 1A) is derived from the previously described pTT vector [37]. The signal peptide sequence, cysteine residues involved in intramolecular cystine formation, and

a)



b)

M<sub>23</sub>ALTFALLVALLVLSCKSSCSVGC<sub>1</sub>DLPQTHSLGSR  
 RTLMLLAQMRRISLFSC<sub>29</sub>LKDRHDFGFPQEEFGNQF  
 QKAETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKF  
 YTELYQQ<sub>98</sub>NDLEAC<sub>98</sub>VIQGVGV<sub>106</sub>TETPLMKEDSILA  
 VRKYFQRITLYLKEKKYSPC<sub>138</sub>AWEVVRAEIMRSFSL  
 STNLQESLRSKE<sub>165</sub>

**Figure 1**

**Expression plasmid encoding human IFN $\alpha$ 2b cDNA. A)** The pYD7-IFN $\alpha$ 2b expression plasmid has been used to generate the D9 clone. (Amp) ampicillin, (Blast) blasticidin, (CMV) cytomegalovirus promoter, (enh MLP) adenovirus major late promoter, (IFN $\alpha$ 2b) human codon-optimized sequence for human IFN $\alpha$ 2b gene, (pA) polyadenylation sequence, (pMB1ori) bacterial origin of replication, (Puro) puromycin, (OriP) Epstein-Barr virus origin of replication, (SV40pA) simian virus 40 polyadenylation sequence, (TPL) adenovirus tripartite leader. **B)** Amino acid sequence of human IFN $\alpha$ 2b. Signal peptide is underlined. The two intramolecular disulfide bridges are C<sub>1</sub>-C<sub>98</sub> and C<sub>29</sub>-C<sub>138</sub>. The glycan-linked threonine (Thr<sub>106</sub>) is underscored.

the threonine of the consensus sequence for O-glycosylation of human IFN $\alpha$ 2b are highlighted (Fig. 1B). The calculated molecular weight of the mature core protein (a.a. 24–188) of IFN $\alpha$ 2b is 19,269 Da. In order to generate IFN $\alpha$ 2b-producing cells, HEK293 were transfected with linearized pYD7-IFN $\alpha$ 2b and selected in the presence of blasticidin. The D9 clone, which stably produces IFN $\alpha$ 2b was isolated as described in material and methods. The production of IFN $\alpha$ 2b with the D9 clone was performed in fed-batch culture. Daily samples from the culture media taken over a period of 9 days were analysed by Coomassie blue-stained gel (Fig. 2A). The gel shows that cell-derived contaminating proteins begin to accumulate significantly after day 5. A decline in cell viability was also notable after day 7 (Fig. 2B). Fed-batch cultivation was terminated and the culture medium harvested. It is noteworthy that early during production, HEK293-derived IFN $\alpha$ 2b migrates with an apparent molecular weight of 2 kDa greater than its predicted mass calculated from the amino acid sequence (19,3 kDa), while at around day 4, a less abundant band of  $\sim$ 19,5 kDa appears. In order to ascertain that this band is also IFN $\alpha$ 2b, N-terminal sequencing was performed on both products. The sequences obtained were identical and read NH<sub>2</sub>-C-D-L-P-Q-T, as expected for N-terminal sequence of human IFN $\alpha$ 2b having a correctly processed signal peptide, therefore suggesting that heterogeneous posttranslational modifications may account for differences in electrophoretic mobilities of these two IFN $\alpha$ 2b species.

#### **Purification of recombinant IFN $\alpha$ 2b by cation exchange chromatography and analysis by gel filtration and SDS-PAGE**

At the end of the production phase, the IFN $\alpha$ 2b is purified as described in the Methods section. The IFN $\alpha$ 2b eluted in a single peak at pH 4,5–4,6 from the cation exchange column (Fig. 3A). The electrophoretic profiles of proteins contained in the harvest, the acid precipitate, the clarified harvest and eluted fractions, are shown on a Coomassie blue-stained gel (Fig. 3B). The acidification step was selective in removing protein contaminants, as the concentration of IFN $\alpha$ 2b in the clarified media was greater than 95% of that quantified in the harvest. The absence of IFN $\alpha$ 2b in the flow through and in the wash suggests that IFN $\alpha$ 2b strongly binds to the SO<sub>3</sub><sup>-</sup> column. According to a conservative estimate performed by densitometric analysis of the SDS-PAGE resolved purified material, the purity of IFN $\alpha$ 2b exceeds 98% after the SO<sub>3</sub><sup>-</sup> column and the final desalting step.

Following desalting in PBS, purified IFN $\alpha$ 2b was loaded on a Superdex 75 gel filtration column. The protein eluted as a single peak with elution volume identical to that of ovalbumin, a 44 kDa protein, indicating that purified HEK293-IFN $\alpha$ 2b is not aggregated and suggesting that it


may form dimers at neutral pH (Fig. 4A). A Coomassie blue-stained gel of IFN-containing fractions shows diffuse wide bands as observed with non-purified material (Fig. 4B). These species with different electrophoretic mobilities reflect glycosylation heterogeneity, which was later confirmed by mass spectroscopy and glycosylation analysis. Under reducing conditions, purified IFN $\alpha$ 2b migrates as a major band of approximately 21 kDa and a less abundant band of lower molecular weight. Mass spectroscopy analysis indicates variations in the molecular weight between 19,922 and 20,659 Da (Fig. 6B). Under non-reducing conditions, IFN $\alpha$ 2b migrates with an apparent molecular weight of  $\sim$ 17 kDa, a greater electrophoretic mobility typical of the presence of intramolecular disulfide bridges (Fig. 4C). The absence of dimers (i.e.  $\sim$ 42 kDa band) in non-reducing conditions indicates that the probable formation of dimers as suggested by gel filtration analysis is independent of intermolecular disulfide bridges. This non-covalent dimeric form of interferon has already been described and involves coordination of a zinc ion by two adjacent glutamic acid residues (E<sub>41</sub> and E<sub>42</sub>) from each of two IFN molecules [38].

#### **The D9 clone produces hundreds of milligrams of IFN $\alpha$ 2b per liter of culture that are efficiently recovered**


IFN $\alpha$ 2b in the crude harvests of fed-batch cultures was quantified by ELISA. The average concentration from two independent productions is  $237 \pm 11$  mg/L, and this was increased to  $301 \pm 25$  mg/L when glucose and glutamine feeds were added during production (Table 1). IFN $\alpha$ 2b recovered from the SO<sub>3</sub><sup>-</sup> column measured by ELISA correlated well with measures obtained with a Bradford assay and by absorbance at 280 nm using IFN $\alpha$ 2b molar extinction coefficient. The concentrations of IFN $\alpha$ 2b measured by ELISA in the harvest and in the recovered fraction from the SO<sub>3</sub><sup>-</sup> column were used to determine the recovery. The mean concentration of IFN $\alpha$ 2b shows that between 70 and 80% of the IFN $\alpha$ 2b produced could be recovered, for two independent productions for each condition (Table 1). These results were comparable in terms of volumetric productivity and recovery to some productions of non-glycosylated IFN $\alpha$ 2b performed in *E. coli* and in the methylotrophic yeast *Pichia pastoris* (Table 2).

#### **IFN $\alpha$ 2b produced in HEK293 is O-glycosylated, highly sialylated and biologically active**


One of the major interests for producing IFN $\alpha$ 2b in mammalian cells is to generate a glycosylated active protein. The apparent molecular weight of IFN $\alpha$ 2b observed on SDS-PAGE suggests that IFN $\alpha$ 2b produced in HEK293 undergoes post-translational modifications. There is also a less abundant product of around 19,5 kDa on SDS-PAGE.

**Figure 2**

**Kinetics of cell growth and IFN $\alpha$ 2b production from D9 clone in fed-batch culture.** D9 cells were seeded at a cell density of  $0.25 \times 10^6$  cells per mL, fed with 0.1% TNI the next day and sampled every day. **A)** Coomassie-stained SDS-PAGE analysis of the culture medium (20  $\mu$ L) collected daily. **B)** Cell counts and viability were measured at the indicated times.

**Figure 3**

**Purification of IFN $\alpha$ 2b by cation-exchange chromatography. A)** A typical chromatographic profile of a purification of HEK293-produced IFN $\alpha$ 2b from a 400 mL fed-batch culture is illustrated. Solid line shows the 280 nm absorbance profile. Dotted line shows pH variations. IFN $\alpha$ 2b elutes in a single peak between 1000 and 1200 mL. **B)** Coomassie-stained SDS-PAGE analysis of 20  $\mu$ L samples collected at different steps of production and purification of IFN $\alpha$ 2b. **1-** crude harvest. **2-** precipitate (equivalent to 200  $\mu$ L of harvest volume). **3-** clarified harvest. **4-** flow through SO $_3^-$  column. **5-** wash SO $_3^-$  column. **6-** elution peak SO $_3^-$  column. **7-** desalted IFN $\alpha$ 2b in PBS.

**Figure 4**

**Purified IFN $\alpha$ 2b is not aggregated and forms dimers at neutral pH independent of intermolecular cystine formation.** Following a desalting step in neutral PBS, purified IFN $\alpha$ 2b was analysed for dimer formation. **A)** Twenty mg of purified IFN $\alpha$ 2b were analysed on a Superdex 75 HR16/60 column equilibrated with PBS at pH 7,0. The arrows and numbers above indicate the elution volumes of molecular weight standards eluted in the same conditions. Purified IFN $\alpha$ 2b elutes in the same volume as ovalbumin, a 44 kDa globular protein. **B)** Coomassie-stained SDS-PAGE analysis of samples (20  $\mu$ L) of each of the 10 fractions (4 mL) collected between elution volumes 40–80 mL. **C)** Coomassie-stained SDS-PAGE analysis of reduced and non-reduced IFN $\alpha$ 2b from HEK293 cells.



**Table 1: Quantification and recovery of HEK293-produced IFN $\alpha$ 2b from two production schemes.**

Production scheme	IFN $\alpha$ 2b mg/L (ELISA)		Percent recovery
	Culture medium	SO <sub>3</sub> -column	
1 feed	237 $\pm$ 11	185 $\pm$ 3	79.5
2 feeds	301 $\pm$ 25	216 $\pm$ 11	71.8

We next determined whether IFN $\alpha$ 2b produced in HEK293 is O-glycosylated [39] and sialylated as previously reported for IFN $\alpha$ 2b produced by human peripheral blood leucocytes [40]. We performed a sequential digestion of purified IFN $\alpha$ 2b with neuraminidase and O-glycosidase to respectively remove sialic acid residues and O-linked saccharides. Each digestion successively increases the electrophoretic mobility of purified IFN $\alpha$ 2b to generate a deglycosylated product that migrates as fast as non-glycosylated recombinant IFN $\alpha$ 2b produced *E. coli* (Fig. 5). This suggests that IFN $\alpha$ 2b produced in HEK293 cells is O-glycosylated and sialylated. Note here the quasi absence of the lower ~19,5 kDa product in the lane containing the non-digested IFN. We found that the majority of this product is lost during the purification process, as most of it remains bound to the column (data not shown). A minor band with lower electrophoretic mobility was still visible after glycosidases treatment, suggesting that this species might be Core 2 type glycan.

A detailed mass analysis and glycosylation pattern of the purified IFN $\alpha$ 2b was next performed by mass spectroscopy. Electrospray ionization (ESI) mass spectrum exhibiting the glycoform profiles associated with each charge state of purified IFN $\alpha$ 2b is shown (Fig. 6A). The masses of


the principal glycoform of this protein correspond to the mature IFN $\alpha$ 2b peptide chain plus the glycans indicated (Fig. 6B). The most intense peak at 20 213 Da appears to be composed of the mature peptide chain plus a single core type-1 disialylated glycan (Hex<sub>1</sub>HexNAc<sub>1</sub>SA<sub>2</sub>). A MS/MS analysis of the tryptic glycopeptides confirms the composition of this glycan. The sialylated (mono and disialylated) glycoforms appear to constitute 75% of the total species. This percentage is likely to be underestimated, as some of the other peaks that cannot be assigned easily may be sialylated as well. The disialylated type 1 glycoform represents 50% of the total peak area while the monosialylated glycoform is 10% of the total. Using electron transfer dissociation, we also show that the glycan is linked to the expected threonine residue at position 106 (Fig. 7).

Finally, we tested the purified glycosylated IFN $\alpha$ 2b produced in HEK293 for *in vitro* biological activity in comparison to non-glycosylated form produced in *E. coli*. Using a reporter gene assay we show that HEK-produced IFN $\alpha$ 2b is biologically active as it induces the production of a secreted alkaline phosphatase (SEAP) reporter enzyme under the control of the human ISG56 promoter (Fig. 8). This assay shows that HEK-produced IFN $\alpha$ 2b is as active

**Table 2: Overview of human recombinant IFN $\alpha$ 2b production in prokaryotic and eukaryotic systems.**

Host	mg/L	Recovery		Purity %	Glycosylation	Activity IU/mg*	Ref.
		mg/L	%				
Prokaryotic							
<i>E. coli</i>	5200	3000	58	ND	No	3 × 10 <sup>9</sup>	[21]
<i>E. coli</i>	4000	300	7,5	ND	No	2,5 × 10 <sup>8</sup>	[19]
<i>E. coli</i>	3500	600	12	100	No	ND	[20]
<i>S. lividans</i>	0,01	ND	ND	ND	No	0,4 × 10 <sup>4</sup>	[55]
Eukaryotic							
<i>Pichia pastoris</i>	450	298	66,2	> 95	ND	1,9 × 10 <sup>9</sup>	[56]
<i>Pichia pastoris</i>	200	ND	ND	ND	No	3.0 × 10 <sup>8</sup>	[57]
Tobacco BY2 cells	0.02	ND	ND	ND	No	ND	[58]
Insect Sf9 cells	ND	ND	ND	ND	Partial (no sialylation)	2,3 × 10 <sup>8</sup>	[34]
Mouse NS0 cells	120	ND	ND	ND		Yes	2 × 10 <sup>8</sup>

\* IFN $\alpha$ 2b activity has been determined by inhibition of viral replication. Different viruses and hosts were used.



**Figure 5**  
**HEK293-produced human IFN $\alpha$ 2b is sialylated and O-glycosylated.** IFN $\alpha$ 2b was deglycosylated as described in material and methods. **1-** 10  $\mu$ g of purified HEK-produced IFN $\alpha$ 2b. **2-** 10  $\mu$ g of purified HEK-produced IFN $\alpha$ 2b digested with neuraminidase. **3-** 10  $\mu$ g of purified HEK-produced IFN $\alpha$ 2b digested with O-glycosidase. **4-** 10  $\mu$ g of purified *E. coli*-produced IFN $\alpha$ 2b.

*in vitro* as bacterially produced IFN $\alpha$ 2b. In addition, viral challenges using Vesicular stomatitis virus (VSV) on Madin-Darby Bovine Kidney (MDBK) cells or using Encephalomyocarditis virus (EMCV) on human A549 cells demonstrated very good antiviral activity of purified IFN $\alpha$ 2b with titres ranging from 4.1 to 12.2  $\times 10^8$  IU/mg for two independent batches (Table 3).

## Discussion


We describe here the generation of a HEK293 cell clone (D9) able to stably produce glycosylated human recombinant IFN $\alpha$ 2b for culture periods up to 4 months in the absence of selection. The volumetric production per litre

of serum-free culture can reach more than 300 mg/L, and is the highest volumetric production of IFN $\alpha$ 2b reported for a mammalian system. We have further developed a rapid and reliable method for its efficient recovery and show that HEK-derived IFN $\alpha$ 2b is O-glycosylated, sialylated and biologically active.


Gel filtration analysis of purified IFN $\alpha$ 2b suggests that it may exist as a dimer in PBS at neutral pH. A zinc-dependent dimeric form of hIFN $\alpha$ 2b as already been observed [38] and was also reported for the structurally homologous human IFN $\beta$  [41]. It is believed that interferon is biologically active as a monomer and the biological significance of zinc-mediated dimerization is currently unknown. The fact that our IFN is biologically active suggests that at the low concentration used in the bioactivity assays, it may exist in solution as a monomer.

To date, the production of recombinant IFN $\alpha$ 2b and other cytokines in mammalian systems, particularly the development of clones stably expressing a cytokine of interest, has not been well exploited due to limitations in the volumetric productivity. One of the possible causes maybe that many cytokines exhibit strong anti-proliferative and cytotoxic activities on diverse cell lines [42,43], therefore strongly selecting against clones that show high cytokine expression levels. The D9 clone nonetheless grew almost as well as parental cells indicating that HEK293 cells can adapt to proliferate in the presence of high levels of IFN $\alpha$ 2b. This adaptability of HEK293 cells to a growth inhibitory cytokine suggests that they may be suitable for the large-scale production of other interferons and cytokines.

To our knowledge, no comparable expression system exists in order to contrast the volumetric productivity of our HEK293 clone. However, this clone performs very well compared to other reported eukaryotic expression systems (see table 2 for an overview of IFN $\alpha$ 2b expression systems). Nevertheless, a much greater volumetric production can be obtained from *E. coli* expression systems. Although we believe that the production capacity of HEK293 cells for IFN $\alpha$ 2b can be improved, we doubt that such productivity can ever be achieved in mammalian cells, at least for a cytokine. It is obvious however that the difficulty in obtaining high recovery of refolded IFN $\alpha$ 2b is still an important challenge with *E. coli*. In general, purifications of recombinant proteins from prokaryotes usually require extraction from inclusion bodies and complex refolding procedures, which reduce recovery yields [44]. Protein refolding is a critical step in the processing of biotherapeutics, as incompletely refolded species lower specific activity and may trigger an immune response. Antibodies to recombinant prokaryotic IFN $\alpha$ 2b have been detected in HCV patients with acquired resistance to

**Figure 6**

**ESI-MS analysis of the intact IFN $\alpha$ 2b glycoprotein. A)** ESI mass spectrum exhibiting the glycoform profiles associated with each charge state of the protein and **B)** the glycoprotein molecule weight profile reconstructed from the mass spectrum in panel A. The most intense peak at 20,213 Da appears to be composed of the mature peptide chain plus a single core type-I disialylated glycan ( $\text{Hex}_1\text{HexNAc}_1\text{SA}_2$ ).

**Figure 7**

**CID and ETD analysis of the tryptic glycopeptides from IFN $\alpha$ 2b. A)** CID-MS/MS spectrum of the triply protonated ion at  $m/z$  1426.8 corresponding to the disialylated glycopeptide of T84-I12. The spectrum is dominated by the sequential neutral loss of the glycan components from the doubly protonated glycopeptide ion. The principal b and y fragment ions arising from fragmentation of the peptide backbone are indicated in the spectrum as are the compositions the glycan oxonium ions observed  $m/z$  494.9, 657.0 and 948.0, respectively. The sequence of the peptide is provided in the inset. **B)** CID-MS/MS spectrum of the triply protonated ion at  $m/z$  1340.8 corresponding to the monosialylated glycopeptide of T84-I12. Note that the neutral loss corresponding to a second sialic acid is missing from this spectrum as is the corresponding oxonium ion at  $m/z$  948.0. **C)** ETD MS/MS spectrum of the triply protonated, monosialylated T84-I12 glycopeptide at  $m/z$  1340.8. The higher  $m/z$  region of the ETD spectrum contained the most informative fragment ions and is presented here. The c ion series indicated in the spectrum clearly identified the site of O-linkage as Threonine 106 of the mature protein.

**Table 3: Assessment of biological activity of HEK293-produced IFN $\alpha$ 2b using two antiviral assays**


Antiviral assay	IFN $\alpha$ 2b activity (U/mg)			
	HEK293 (1 feed)	<i>E. coli</i>	HEK293 (2 feeds)	<i>E. coli</i>
MDBK/VSV	$4.1 \times 10^8$	$3.1 \times 10^8$	$12.2 \times 10^8$	$4.0 \times 10^8$
A549/EMCV	$6.0 \times 10^8$	$3.9 \times 10^8$	$7.2 \times 10^8$	$8.1 \times 10^8$

IFN $\alpha$ 2b activity has been determined by inhibition of viral replication of vesicular stomatitis virus (VSV) in Madin Darby bovine kidney cells (MDBK) and of encephalomyocarditis virus (EMCV) in the lung carcinoma cell line A-549. Both antiviral assays were carried out by PBL InterferonSource. In each assay performed independently, *E. coli* IFN $\alpha$ 2b was used as a positive control.

IFN $\alpha$ 2b treatment [26,27], although it is not clear whether denatured IFN $\alpha$ 2b played a role in this case.

Because the vast majority of biotherapeutics including growth factors, cytokines and antibodies are secreted proteins, mammalian systems, unlike prokaryotes, allow for production in perfusion as well as for the development of non-denaturing purification procedures. The first and foremost advantage of producing human recombinant proteins in mammalian systems is to generate proteins with the necessary posttranslational modifications required for full biological activity. N-glycosylation in particular, is often required for proper protein folding [45], protein-protein interactions, stability and optimal pharmacokinetics [46]. Although O-glycosylation is less

critical for structure and function of proteins, it has been shown for example to increase the serum half-life of IGFBP6 by 2,3 folds over the non-glycosylated protein [47] and to protect against proteolysis [48]. In a recent randomized study, O-glycosylated IFN $\alpha$ 2b was shown to have an increased serum half life in comparison to non-glycosylated IFN $\alpha$ 2b [32]. We show here that human recombinant IFN $\alpha$ 2b produced in HEK293 cells is O-glycosylated and extensively sialylated. Despite heterogeneity in the glycan structures, the nature and distribution of glycan moieties are quite similar to IFN $\alpha$ 2b naturally produced by human leukocytes [40]. Approximately 50% of the purified protein is disialylated, while another 10% is monosialylated, in comparison to 50% and 30% respectively for leukocyte-derived IFN. As expected from its biochemical structure, we show that HEK293-produced IFN $\alpha$ 2b has a biological activity comparable to that of non-glycosylated *E. coli*-produced IFN $\alpha$ 2b by means of an *in vitro* reporter-gene assay, indicating that IFN $\alpha$ 2b is not inactivated by the purification process. This was also confirmed by the high specific activity ranging from 4.1 to  $12.2 \times 10^8$  IU/mg using two different batches of purified IFN and two antiviral assays.



**Figure 8**  
**HEK293-produced human IFN $\alpha$ 2b is biologically active.** The biological activity of HEK293-produced human IFN $\alpha$ 2b was assayed with a gene reporter assay and compared to *E. coli*-produced human recombinant IFN $\alpha$ 2b as described in material and methods. The activity of the secreted alkaline phosphatase is plotted against the concentration of IFN $\alpha$ 2b produced in the two hosts. Each point represents the average  $\pm$  SEM of 3 experiments performed in triplicate.

## Conclusion

While additional studies are needed to determine whether HEK293-produced IFN $\alpha$ 2b can offer advantages over non-glycosylated or pegylated IFN for *in vivo* applications, this work demonstrates that the HEK293 cell line is a suitable host for the high volumetric production of glycosylated human recombinant IFN $\alpha$ 2b and potentially other cytokines.

## Methods

### Material

The expression plasmid was purified with a maxi-prep plasmid purification kit (Qiagen, Mississauga, ON, Canada). F17 serum-free culture media and blasticidin were obtained from Invitrogen (Carlsbad, CA). Pluronic F68 and glutamine were from Sigma-Aldrich (St. Louis, MO) and Tryptone N1 from Organotechnie (La Courneuve, France). Reagents for IFN $\alpha$ 2b purification and electrophoresis include anhydrous citric acid and tri-Na citrate

(EMD Chemicals Inc, Darmstadt, Germany), 0.45  $\mu$ m filtering units (Millipore, Bedford, MA), NaCl (Sigma-Aldrich, St. Louis, MO), Fractogel<sup>®</sup> SO3<sup>-</sup> (M) (Merck KGaA, Darmstadt, Germany), Econo-Pac<sup>®</sup> 10 columns (Bio-Rad Laboratories), Bradford Reagent (Biorad, Hercules, CA), 2  $\mu$ m filters (Pall Corp, Ann Arbor, MI), NuPAGE Bis Tris 4–12% gradient gels, MES 20 $\times$  buffer (Invitrogen, Carlsbad, CA), and Coomassie R250 stain (Sigma-Aldrich, St. Louis, MO). Trypsin (Promega, Madison WI), neuraminidase, dithiothreitol, iodoacetamide and guanidine HCl (Sigma-Aldrich, St. Louis, MO), O-glycosidase (Roche), Tris HCl, (Bio-Rad, Mississauga, ON), high purity acetonitrile, formic acid and ammonium bicarbonate (VWR International, Montreal, QC) and Centricon 3,000 MWL centrifugal filters (Millipore, Bedford, MA) were used for glycosylation analysis. The IFN $\alpha$  antibody and ELISA kit are from PBL InterferonSource (Piscataway, NJ, USA) and bacterially produced IFN $\alpha$ 2b from Cell Sciences Inc (Norwood, MA, USA). pNifty2-56K-SEAP plasmid is from Invivogen (San Diego, USA).

#### **IFN $\alpha$ 2b expression plasmid**

The IFN $\alpha$ 2b gene was synthesized with human-optimized codons (Geneart AG, Regensburg, Germany) according to the Genebank sequence no. AY255838. The synthetic cDNA was inserted as a BamHI/EcoRI fragment downstream of the cytomegalovirus (CMV) promoter into the pYD7 expression plasmid. This plasmid is a derivative of the previously described pTT vector [37] encoding the original functional elements in addition to a blasticidin resistance cassette.

#### **Engineering of a HEK293 clone stably expressing IFN $\alpha$ 2b and fed-batch production**

A HEK293 cell line constitutively expressing the EBNA1 protein of EBV (clone 6E) was used to generate IFN-producing clones. HEK293-6E and IFN-producing clone are grown in suspension in serum-free F17 culture media supplemented with 0.1% pluronic F68. Cultures were grown at 37°C and 5% CO<sub>2</sub> under constant agitation (120 rpm). HEK293-6E were transfected as previously described [37] with PvuI-linearized pYD7/IFN $\alpha$ 2b and selected in the presence of 2  $\mu$ g/mL of blasticidin. The blasticidin resistant cells were next seeded into 96 well plates at 1 cell/well without blasticidin. After 3–4 weeks, emerging clones were expanded (in the absence of blasticidin) and tested for IFN $\alpha$ 2b expression by dot blot. The selection of IFN-producing clones was based on the levels of IFN $\alpha$ 2b expression and growth properties of the clones. The highest producers were amplified as suspension cultures and tested for IFN $\alpha$ 2b accumulation over a 4 days culture. One clone, identified as D9, was selected because it is stably producing high IFN $\alpha$ 2b levels while maintaining a high growth rate (doubling time of 26 hours<sup>-1</sup>). For IFN $\alpha$ 2b production, cells were seeded at a density of 0,25

$\times 10^6$  cells/mL in F17 antibiotic-free media in shaker flasks. Twenty-four hours post-seeding, the cultures were fed with 0,5% peptones [37,49] and left in the incubator for an additional 7–8 days. Optional addition of 20 mM glucose and 5 mM glutamine was performed 4 days post seeding where indicated.

#### **Purification of IFN $\alpha$ 2b**

The culture medium of a fed-batch culture was collected by centrifugation at 1000 g for 10 min. The supernatant was then acidified to pH 3.6–3.8 with 1 M citric acid. Acidification caused the formation of a precipitate which was removed by centrifugation. The clarified supernatant was then filtered on a 0.45  $\mu$ m filtering unit. Purification of IFN $\alpha$ 2b from the filtered supernatant was performed on an ÄKTA Explorer system (GE healthcare, Baie D'Urfé, QC, Canada). The supernatant was loaded at a flow rate of 10 mL/min on a Fractogel SO3-cation exchange column, previously equilibrated with 0,1 M Tri-Na citrate buffer pH 3,5 containing 0,35 M NaCl. Following a wash with 2 column volumes of the equilibration buffer, the IFN $\alpha$ 2b was then eluted with a pH gradient. The pH of the mobile phase was increased from pH 3,5 to pH 6,0 with 0,1 M Tri-Na citrate buffer pH 6,0, plus 0,35 M NaCl. The fractions containing IFN $\alpha$ 2b were pooled. An additional desalting step was performed on Econo-Pac<sup>®</sup> 10 columns according to the manufacturer's specifications. For the determination of glycosylation by enzymatic digestion, the purified IFN $\alpha$ 2b was desalted in 0.1 M NH<sub>4</sub>HCO<sub>3</sub> buffer pH 5 and lyophilized, whereas for bioassays, the purified IFN $\alpha$ 2b was desalted in PBS and sterile filtered.

#### **Quantification and purity of IFN $\alpha$ 2b**

IFN $\alpha$ 2b recovered from the SO3<sup>-</sup> column was quantified by measuring absorption at 280 in a spectrophotometer, with a Nanodrop ND-1000 (Fisher Scientific, Montreal, QC, Canada), with a Bradford assay and by ELISA according to the manufacturer's protocol. The concentration in the harvest was measured with ELISA and used to calculate the percent recovery. To assess the purity level of IFN $\alpha$ 2b, 3  $\mu$ g were analyzed by SDS-PAGE followed by Coomassie staining.

#### **N-terminal sequencing and enzymatic determination of glycosylation of purified IFN $\alpha$ 2b**

As HEK293-produced IFN $\alpha$ 2b migrates as two bands on SDS-PAGE, N-terminal amino acid sequences from both bands were obtained by automated sequencing performed at our sequencing facility. Enzymatic treatments with neuraminidase and O-glycosidase were performed to remove sialic acid and O-linked sugars respectively. Sequential digestions were performed in 50 mM phosphate buffer pH 5,0 on 100  $\mu$ g of purified/desalted IFN $\alpha$ 2b. Removal of sialic acid was done with 0.5 IU of neuraminidase for 1 h at 37°C followed by addition of 15

mU of O-glycosidase. Non-glycosylated recombinant IFN $\alpha$ 2b from *E. coli* and glycosylated and deglycosylated HEK293-produced IFN $\alpha$ 2b were resolved on SDS-PAGE in parallel to compare migration profiles. Migration profiles of glycosylated and deglycosylated HEK293-produced IFN $\alpha$ 2b were compared to non-glycosylated IFN $\alpha$ 2b produced in *E. coli*.

#### **Analysis of intact IFN $\alpha$ 2b by mass spectrometry**

The protein solution ( $\sim 1 \mu\text{g}/\mu\text{L}$  in PBS buffer) was desalted by filtration on a 3 000 MWL Centricon filter and diluted to its original concentration with deionized water. The solution was adjusted to 20% acetonitrile, 0.2% formic acid just prior to infusion at  $1 \mu\text{L}/\text{min}$  into the electrospray interface of a Q-TOF 2 hybrid quadrupole time-of-flight mass spectrometer (Waters, Milford, MA). The mass spectrometer was set to acquire one spectrum every 2 seconds over the mass range,  $m/z$  800–2600. The protein molecular weight profile was generated from the mass spectrum using MaxEnt (Waters).

#### **Sequence analysis of the tryptic glycopeptides from purified IFN $\alpha$ 2b**

Purified IFN $\alpha$ 2b was reduced, alkylated and digested with trypsin according to standard protocols. In summary, approximately 100  $\mu\text{g}$  of the protein was dissolved in 1M Tris HCl, 6M guanidine HCl, pH 7.5 containing 2 mM dithiothreitol (DTT) and incubated at 50°C for 1 hour. The reduced cysteines were converted to carboxyamido-methyl derivatives using 10-fold excess of iodoacetamide over DTT. The protein solution was then concentrated on a 3 000 MWL Centricon and diluted to 100  $\mu\text{L}$  using 50 mM ammonium bicarbonate. This process was repeated a second time. Trypsin (5  $\mu\text{g}$ ) was added to the sample, which was then incubated overnight at 37°C.

The tryptic digest was analyzed and fractionated by LC-MS using an Agilent 1100 HPLC system coupled with the Q-TOF2 mass spectrometer. Approximately 60  $\mu\text{g}$  of the protein digest was injected onto a 4.6 mm  $\times$  250 cm Jupiter, 5  $\mu\text{m}$ , 300 Å, C18 column (Phenomenex, Torrance, CA) and resolved using the following gradient conditions: 5% to 60% acetonitrile, 0.2% formic acid in 45 minutes, increasing to 95% after 50 minutes (1 mL/min flow rate). Approximately, 60  $\mu\text{L}/\text{min}$  of the HPLC eluate was directed to the mass spectrometer while the remainder was collected in 1 minute fractions. The Q-TOF2 mass spectrometer was set to acquire 1 spectrum per second ( $m/z$  150–2000) whilst cycling between a low and high offset voltage within the collision cell (10 V and 35V, respectively). This enabled the simultaneous detection of intact peptide and glycopeptide ions in the higher  $m/z$  regions (low offset mode) as well as the unique glycan oxonium ions in the lower regions of the spectrum (high

offset mode). By screening the fractions in this manner it was possible to determine that only two of them (fractions 25–26 and 26–27 minutes, respectively) contained glycopeptides.

Glycopeptides were interrogated by collision induced dissociation (CID) to determine their amino acid sequence and glycan composition and by electron transfer dissociation (ETD) to identify the site of linkage. ETD preserves delicate modifications intact during the fragmentation process and is ideal for identifying the linkage sites of O-glycans [50–52]. The glycopeptide-containing fractions were infused at  $1 \mu\text{L}/\text{min}$  into the electrospray ionization source of a LTQ XL linear ion trap (Thermo Fisher Scientific) capable of performing ETD. The CID collision voltage was adjusted for optimum production of peptide fragment ions from the multiply charge glycopeptide precursor ions (typically 25–30 V). ETD was performed using fluoranthene as the anionic reagent and with supplementary activation enabled. The optimal ETD reaction time for these glycopeptides was 350 msec.

#### **Biological activity**

A SEAP reporter gene assay based on expression plasmid containing an IFN-inducible promoter (pNiFty2) was used to assess the biological activity of glycosylated HEK293-produced IFN $\alpha$ 2b in comparison to non-glycosylated IFN $\alpha$ 2b. HEK293 cells were transfected with the pNiFty2 reporter plasmid, which encodes the secreted embryonic alkaline phosphatase (SEAP) under the control of the human ISG56 promoter. Transfected cells were plated in 96 well plates at a cell density of  $10^5$  cells/mL and stimulated, 24 h post-transfection, with IFN $\alpha$ 2b at the indicated concentrations. Following an additional 48 h period of incubation, the supernatants were collected and assayed for SEAP activity. The hydrolysis of paranitrophenyl phosphate (pNPP) was measured as a function of time to determine SEAP activity induced with IFN treatments, as previously described [53]. The SEAP activity is expressed as the increase in absorbance units at 410 nm per minute.

Antiviral assays were carried out by PBL InterferonSource (Piscataway, NJ) using Madin-Darby Bovine Kidney (MDBK) cells challenged by Vesicular stomatitis virus or human A549 cells challenged with encephalomyocarditis virus [54]. Cells were incubated with two-fold serial dilutions of IFN $\alpha$ 2b standard (PBL InterferonSource), 293-IFN $\alpha$ 2b, or control (Media). After 24 h incubation with the virus, cell viability was determined via crystal violet staining (570 nm absorbance). The sample titer (IC<sub>50</sub>), calculated by SigmaPlot software (SPSS Inc., Point Richmond, CA), was based on the 50% cytopathic effect of the assay. Units of anti-viral activity were based on the titer of a PBL lab standard, which was determined against the

NIH reference standard for human IFN $\alpha$ 2b (Gxa01-901-535).

## Abbreviations

EBV: Epstein Barr Virus; ESI: electrospray ionization; ETD: electron transfer dissociation; HEK: human embryonic kidney; IFN $\alpha$ 2b: interferon alpha2b; PBS: phosphate buffered saline; SEAP: secreted alkaline phosphatase.

## Authors' contributions

ML performed production of IFN $\alpha$ 2b, determined cell viability and growth curves in fed-batch productions, co-developed the purification method and purified IFN $\alpha$ 2b, performed enzymatic deglycosylation, interpreted the data and wrote the manuscript. SP did ELISA assays and characterized the D9 IFN-producing clone. JK contributed the analysis of glycosylation by mass spectroscopy and editing of the manuscript. LB performed batch and fed-batch developments. DB performed the transfection and isolation of the D9 clone. BC co-developed the purification method and performed some batch experiments. FA tested IFN $\alpha$ 2b for bioactivity. YD designed experiments, interpreted the data and revised the manuscript.

## Acknowledgements

We thank France Dumas for N-terminal sequencing of IFN $\alpha$ 2b, Alain Drouin, Wen Ding and Luc Tessier for their technical assistance with the mass spectrometry analysis and Phuong Lan Pham for insightful discussion.

## References

- Chelbi-Alix MK, Wietzerbin J: **Interferon, a growing cytokine family: 50 years of interferon research.** *Biochimie* 2007, **89**:713-718.
- Kaluz S, Kabat P, Gibadulinova A, Vojtassak J, Fuchsberger N, Kontsek P: **Interferon alpha2b is the predominant subvariant detected in human genomic DNAs.** *Acta Virol* 1994, **38**:101-104.
- Gewert DR, Sharp NA, Barber KA, Cooper H, Tucker D, Lewis AP, Thursz M, Crowe JS: **Detection of rare allelic variants of the interferon-alpha 2 gene in human genomic DNA.** *J Interferon Cytokine Res* 1995, **15**:403-406.
- Mohanty SR, Kupfer SS, Khiani V: **Treatment of chronic hepatitis B.** *Nat Clin Pract Gastroenterol Hepatol* 2006, **3**:446-458.
- Vogel W: **Treatment of acute hepatitis C virus infection.** *J Hepatol* 1999, **31 Suppl 1**:189-192.
- Bajetta E, Del VM, Nova P, Fusi A, Daponte A, Sertoli MR, Queirolo P, Taveggia P, Bernengo MG, Legha SS, Formisano B, Cascinelli N: **Multicenter phase III randomized trial of polychemotherapy (CVD regimen) versus the same chemotherapy (CT) plus subcutaneous interleukin-2 and interferon-alpha2b in metastatic melanoma.** *Ann Oncol* 2006, **17**:571-577.
- Terheyden P, Becker JC, Kampgen E, Brocker EB: **Sequential interferon-alpha2b, interleukin-2 and fotemustine for patients with metastatic melanoma.** *Melanoma Res* 2000, **10**:475-482.
- Eggermont AM, Suciu S, MacKie R, Ruka W, Testori A, Kruit W, Punt CJ, Delauney M, Sales F, Groenewegen G, Ruiter DJ, Jagiello I, Stoitchkov K, Keilholz U, Lienard D: **Post-surgery adjuvant therapy with intermediate doses of interferon alpha 2b versus observation in patients with stage IIb/III melanoma (EORTC 18952): randomised controlled trial.** *Lancet* 2005, **366**:1189-1196.
- Dezube BJ: **New therapies for the treatment of AIDS-related Kaposi sarcoma.** *Curr Opin Oncol* 2000, **12**:445-449.
- Angstreich GR, Smith BD, Jones RJ: **Treatment options for chronic myeloid leukemia: imatinib versus interferon versus allogeneic transplant.** *Curr Opin Oncol* 2004, **16**:95-99.
- Bonifazi F, de VA, Rosti G, Guilhot F, Guilhot J, Trabacchi E, Hehlmann R, Hochhaus A, Shepherd PC, Steegmann JL, Kluin-Nelemans HC, Thaler J, Simonsson B, Louwagie A, Reiffers J, Mahon FX, Montefusco E, Alimena G, Hasford J, Richards S, Saglio G, Testoni N, Martinelli G, Tura S, Baccarani M: **Chronic myeloid leukemia and interferon-alpha: a study of complete cytogenetic responders.** *Blood* 2001, **98**:3074-3081.
- Marler JJ, Rubin JB, Trede NS, Connors S, Grier H, Upton J, Mulliken JB, Folkman J: **Successful antiangiogenic therapy of giant cell angioblastoma with interferon alpha 2b: report of 2 cases.** *Pediatrics* 2002, **109**:E37.
- Pearlman BL: **Hepatitis C treatment update.** *Am J Med* 2004, **117**:344-352.
- Manns MP, McHutchison JG, Gordon SC, Rustgi VK, Shiffman M, Rein-dollar R, Goodman ZD, Koury K, Ling M, Albrecht JK: **Peginterferon alpha-2b plus ribavirin compared with interferon alpha-2b plus ribavirin for initial treatment of chronic hepatitis C: a randomised trial.** *Lancet* 2001, **358**:958-965.
- Hoofnagle JH, Seeff LB: **Peginterferon and ribavirin for chronic hepatitis C.** *N Engl J Med* 2006, **355**:2444-2451.
- Allain JP: **Epidemiology of Hepatitis B virus and genotype.** *J Clin Virol* 2006, **36 Suppl 1**:S12-S17.
- Seeff LB, Hoofnagle JH: **Epidemiology of hepatocellular carcinoma in areas of low hepatitis B and hepatitis C endemicity.** *Oncogene* 2006, **25**:3771-3777.
- Picozzi VJ, Kozarek RA, Traverso LW: **Interferon-based adjuvant chemoradiation therapy after pancreaticoduodenectomy for pancreatic adenocarcinoma.** *Am J Surg* 2003, **185**:476-480.
- Babu KR, Swaminathan S, Marten S, Khanna N, Rinas U: **Production of interferon-alpha in high cell density cultures of recombinant Escherichia coli and its single step purification from refolded inclusion body proteins.** *Appl Microbiol Biotechnol* 2000, **53**:655-660.
- Beldarrain A, Cruz Y, Cruz O, Navarro M, Gil M: **Purification and conformational properties of a human interferon alpha2b produced in Escherichia coli.** *Biotechnol Appl Biochem* 2001, **33**:173-182.
- Srivastava P, Bhattacharaya P, Pandey G, Mukherjee KJ: **Overexpression and purification of recombinant human interferon alpha2b in Escherichia coli.** *Protein Expr Purif* 2005, **41**:313-322.
- Youngster S, Wang YS, Grace M, Bausch J, Bordens R, Wyss DF: **Structure, biology, and therapeutic implications of pegylated interferon alpha-2b.** *Curr Pharm Des* 2002, **8**:2139-2157.
- Glue P, Fang JW, Rouzier-Panis R, Raffanel C, Sabo R, Gupta SK, Salfi M, Jacobs S: **Pegylated interferon-alpha2b: pharmacokinetics, pharmacodynamics, safety, and preliminary efficacy data.** *Hepatitis C Intervention Therapy Group.* *Clin Pharmacol Ther* 2000, **68**:556-567.
- Vyas K, Brassard DL, Delorenzo MM, Sun Y, Grace MJ, Borden EC, Leaman DW: **Biologic activity of polyethylene glycol 2000-interferon-alpha2b compared with interferon-alpha2b: gene modulatory and antigrowth effects in tumor cells.** *J Immunother* 2003, **26**:202-211.
- Grace MJ, Lee S, Bradshaw S, Chapman J, Spond J, Cox S, DeLorenzo M, Brassard D, Wylie D, Cannon-Carlson S, Cullen C, Indelicato S, Voloch M, Bordens R: **Site of pegylation and polyethylene glycol molecule size attenuate interferon-alpha antiviral and antiproliferative activities through the JAK/STAT signaling pathway.** *J Biol Chem* 2005, **280**:6327-6336.
- Giannelli G, Antonelli G, Fera G, Del VS, Riva E, Broccia C, Schiraldi O, Dianzani F: **Biological and clinical significance of neutralizing and binding antibodies to interferon-alpha (IFN-alpha) during therapy for chronic hepatitis C.** *Clin Exp Immunol* 1994, **97**:4-9.
- van der Eijk AA, Vrolijk JM, Haagsmans BL: **Antibodies neutralizing peginterferon alpha during retreatment of hepatitis C.** *N Engl J Med* 2006, **354**:1323-1324.
- Braun A, Kwee L, Labow MA, Alsenz J: **Protein aggregates seem to play a key role among the parameters influencing the antigenicity of interferon alpha (IFN-alpha) in normal and transgenic mice.** *Pharm Res* 1997, **14**:1472-1478.
- Bianchi AA, McGrew JT: **High-level expression of full-length antibodies using trans-complementing expression vectors.** *Biotechnol Bioeng* 2003, **84**:439-444.
- Grunberg J, Knogler K, Waibel R, Novak-Hofer I: **High-yield production of recombinant antibody fragments in HEK-293 cells using sodium butyrate.** *Biotechniques* 2003, **34**:968-972.



31. Wurm FM: **Production of recombinant protein therapeutics in cultivated mammalian cells.** *Nat Biotechnol* 2004, **22**:1393-1398.
32. Patel TB, Pequignot E, Parker SH, Leavitt MC, Greenberg HE, Kraft WK: **Transgenic avian-derived recombinant human interferon-alpha2b (AVI-005) in healthy subjects: an open-label, single-dose, controlled study.** *Int J Clin Pharmacol Ther* 2007, **45**:161-168.
33. Rapp JC, Harvey AJ, Speksnijder GL, Hu W, Ivarie R: **Biologically active human interferon alpha-2b produced in the egg white of transgenic hens.** *Transgenic Res* 2003, **12**:569-575.
34. Sugiyama K, Ahorn H, Maurer-Fogy I, Voss T: **Expression of human interferon-alpha 2 in Sf9 cells. Characterization of O-linked glycosylation and protein heterogeneities.** *Eur J Biochem* 1993, **217**:921-927.
35. Zwarthoff EC, Bosveld IJ, Vonk WP, Trapman J: **Constitutive expression of a murine interferon alpha gene in hamster cells and characterization of its protein product.** *J Gen Virol* 1985, **66** (Pt 4):685-691.
36. Rossmann C, Sharp N, Allen G, Gewert D: **Expression and purification of recombinant, glycosylated human interferon alpha 2b in murine myeloma NSo cells.** *Protein Expr Purif* 1996, **7**:335-342.
37. Durocher Y, Perret S, Kamen A: **High-level and high-throughput recombinant protein production by transient transfection of suspension-growing human 293-EBNA1 cells.** *Nucleic Acids Res* 2002, **30**:E9.
38. Radhakrishnan R, Walter LJ, Hruza A, Reichert P, Trotta PP, Nagabhushan TL, Walter MR: **Zinc mediated dimer of human interferon-alpha 2b revealed by X-ray crystallography.** *Structure* 1996, **4**:1453-1463.
39. Adolf GR, Kalsner I, Ahorn H, Maurer-Fogy I, Cantell K: **Natural human interferon-alpha 2 is O-glycosylated.** *Biochem J* 1991, **276** (Pt 2):511-518.
40. Nyman TA, Kalkkinen N, Tolo H, Helin J: **Structural characterisation of N-linked and O-linked oligosaccharides derived from interferon-alpha2b and interferon-alpha14c produced by Sendai-virus-induced human peripheral blood leukocytes.** *Eur J Biochem* 1998, **253**:485-493.
41. Karpusas M, Nolte M, Benton CB, Meier W, Lipscomb WN, Goelz S: **The crystal structure of human interferon beta at 2.2-A resolution.** *Proc Natl Acad Sci U S A* 1997, **94**:11813-11818.
42. Pokrovskaja K, Panaretakis T, Grander D: **Alternative signaling pathways regulating type I interferon-induced apoptosis.** *J Interferon Cytokine Res* 2005, **25**:799-810.
43. van Boxel-Dezaire AH, Rani MR, Stark GR: **Complex modulation of cell type-specific signaling in response to type I interferons.** *Immunity* 2006, **25**:361-372.
44. Baneyx F, Mujacic M: **Recombinant protein folding and misfolding in Escherichia coli.** *Nat Biotechnol* 2004, **22**:1399-1408.
45. Molinari M: **N-glycan structure dictates extension of protein folding or onset of disposal.** *Nat Chem Biol* 2007, **3**:313-320.
46. Werner RG, Kopp K, Schlueter M: **Glycosylation of therapeutic proteins in different production systems.** *Acta Paediatr Suppl* 2007, **96**:17-22.
47. Marinaro JA, Casley DJ, Bach LA: **O-glycosylation delays the clearance of human IGF-binding protein-6 from the circulation.** *Eur J Endocrinol* 2000, **142**:512-516.
48. Marinaro JA, Neumann GM, Russo VC, Leeding KS, Bach LA: **O-glycosylation of insulin-like growth factor (IGF) binding protein-6 maintains high IGF-II binding affinity by decreasing binding to glycosaminoglycans and susceptibility to proteolysis.** *Eur J Biochem* 2000, **267**:5378-5386.
49. Pham PL, Perret S, Cass B, Carpentier E, St-Laurent G, Bisson L, Kamen A, Durocher Y: **Transient gene expression in HEK293 cells: peptone addition posttransfection improves recombinant protein synthesis.** *Biotechnol Bioeng* 2005, **90**:332-344.
50. Coon JJ, Ueberheide B, Syka JE, Dryhurst DD, Ausio J, Shabanowitz J, Hunt DF: **Protein identification using sequential ion/ion reactions and tandem mass spectrometry.** *Proc Natl Acad Sci U S A* 2005, **102**:9463-9468.
51. Hogan JM, Pittner SJ, Chrisman PA, McLuckey SA: **Complementary structural information from a tryptic N-linked glycopeptide via electron transfer ion/ion reactions and collision-induced dissociation.** *J Proteome Res* 2005, **4**:628-632.
52. Syka JE, Coon JJ, Schroeder MJ, Shabanowitz J, Hunt DF: **Peptide and protein sequence analysis by electron transfer dissociation mass spectrometry.** *Proc Natl Acad Sci U S A* 2004, **101**:9528-9533.
53. Durocher Y, Perret S, Thibaudau E, Gaumond MH, Kamen A, Stocco R, Abramovitz M: **A reporter gene assay for high-throughput screening of G-protein-coupled receptors stably or transiently expressed in HEK293 EBNA cells grown in suspension culture.** *Anal Biochem* 2000, **284**:316-326.
54. Rubinstein S, Familletti PC, Pestka S: **Convenient assay for interferons.** *J Virol* 1981, **37**:755-758.
55. Pimienta E, Fando R, Sanchez JC, Vallin C: **Secretion of human interferon alpha 2b by Streptomyces lividans.** *Appl Microbiol Biotechnol* 2002, **58**:189-194.
56. Shi L, Wang D, Chan W, Cheng L: **Efficient expression and purification of human interferon alpha2b in the methylotrophic yeast, Pichia pastoris.** *Protein Expr Purif* 2007, **54**:220-226.
57. Ghosalkar A, Sahai V, Srivastava A: **Secretory expression of interferon-alpha 2b in recombinant Pichia pastoris using three different secretion signals.** *Protein Expr Purif* 2008.
58. Xu J, Tan L, Goodrum KJ, Kieliszewski MJ: **High-yields and extended serum half-life of human interferon alpha2b expressed in tobacco cells as arabinogalactan-protein fusions.** *Biotechnol Bioeng* 2007, **97**:997-1008.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
http://www.biomedcentral.com/info/publishing\_adv.asp

